

Studies on the Quality and Heavy Metals in Common Salt Production Using Sea Brine and Sub-Soil Brine in South Tamil Nadu, India

P. Prema Kumari¹, K. Mercy Salima² and P. Ethalsha³

^{1&2}Humanities and Sciences Department, N. M. S. Kamaraj Polytechnic College, Pazhavilai, Tamil Nadu, India

³Research Scholar, Department of Chemistry, Scott Christian College, Tamil Nadu, India

E-Mail: prema2010phd@gmail.com

(Received 11 April 2015; Revised 30 April 2015; Accepted 18 May 2015; Available online 27 May 2015)

Abstract - The manufacture of common salt using sea brine and sub-soil brine contain several chemical and trace metal elements. This study estimated the salt quality parameters like moisture content, insoluble matter, calcium sulphate, magnesium sulphate, magnesium chloride and sodium chloride content of the salt and also estimated the different trace metals like copper, iron, zinc, manganese and chromium in salts harvested from the Salt samples of two selected salt works, Kovalam SW₁ (sea brine) and Puthalam SW₂ (sub-soil brine) in Kanyakumari District and the two selected salt works, Vepalodai SW₃ (sea brine) and Sahupuram SW₄ (sub-soil brine) in Tuticorin District. The salt samples have maximum moisture content in Kanyakumari salt works than Tuticorin District because of the climatic conditions. The salt works using sea brine, harvested salt of the maximum mean sodium chloride content than sub-soil brine using salt works. In both the Districts, sea brine using salt works is having almost similar concentration of heavy metals, likewise sub soil brine using salt works.

Keywords: Quality, Common Salt Production, Sea Brine, Sub-Soil Brine

I. INTRODUCTION

Sea salt contains 92 essential minerals and most all refined adulterated sea salts contain only 2 elements (Na⁺ and Cl⁻). Biologically, 24 of these elements in real sea salt have already been proven necessary and essential to maintain and recover health (Earl Friden, 1972). The rice eating population requires more salt than others, because rice is deficient in salt. In temperate climate, the annual human requirement of salt is about 5kg/year and it is more in the tropical countries (D.W.Kaufmann, 1960). Sodium and chloride are found in almost all foods. Salt is added to many pet foods, and as we know, can increase the flavour of foods. Sodium and chloride may also be bound to minerals or molecules such as vitamin K or potassium (K). Salt is produced mainly by solar evaporation of sea water, lake and sub soil brines. More than 50% of the countries undergo solar salt production. While the process for solar evaporation of brine is same around the world over, the manufacturing techniques and product quality vary considerably (United Nations Industrial Development Organisation, 1968).

The quality and quantity of the salt depends upon the quality of the brine used, the salt as marketed should not contain more than 6.5% moisture and not more than 1.5 %

insoluble matter. The associated salt i.e., magnesium chloride should not exceed 1.2% and no other allied salt may exceed 1% (S.C.Agarwall, 1956). The technique of solar salt production involves fractional crystallization of salts in different ponds to get common salt in the purest form possible (P.Sorgeloos *et al.*, 1990). Common salt is essentially sodium chloride and its quality is determined by its sodium chloride content, the higher the percentage, the purer the salt. Besides human requirement of the salt of good quality, high purity of salt is necessary for all kinds of heavy chemical industries that use salt as their raw material. But, the impurities in the sea salt are unavoidable. In fact the multiplicity of impurities in salt, their relative quantities and how they influence the salt properties are so variable that every salt needs to be considered on its own merits.

Thus the impurities and the various elements present in source water greatly influence the quality of salt. Except for insolubles, the origin of impurities is the sea water. Calcium sulphate is the most persistent companion of sea salt. Magnesium salts are always present in the sea salt, usually in the ratio of approximately one and a half weight units of magnesium chloride and one weight of magnesium sulphate. In America and other advanced countries, salt of as high as 99.9% purity is manufactured and offered for sale. , salt produced in properly designed salt works has a purity of 90 to 95% NaCl, 1% calcium salts, 1 to 2% magnesium salts and 5 to 8% water. If the salt is washed and dried, its purity can be improved upto 99%. According to the typical standards adapted in the developing countries the purity required is 99.5% for grade I industrial salt, 98.5% for grade II industrial salt, 97% for table salt and 96% for edible common salt. In the developed countries the specifications are more stringent, the minimum purity prescribed for table salt being 99.5% (G.Leemarose, 2007). Thus high quality of salt is expected not only for human consumption but also for various other uses.

The term 'trace metal' may describe a metal found in trace amounts in an organism (eg., less than 0.01%) of the mass of organism or may have a further restriction and apply only to those metals required in metabolism (Rainbow, 1986).

Manganese is required for the enzyme needed to synthesize the mucopoly sacharide in cartilage and bone. It also plays a key role in activating enzymes required for carbohydrate

and lipid metabolism problems associated with manganese deficiency include weakness, slow growth, shift joints and high birth mortality. Manganese compounds appear to be the most effective scavengers of other trace elements (P.Miramand *et al.*, 2001). Along with iron, copper is also necessary in haemoglobin formation. Zinc is an essential micronutrient in all marine organisms, being a cofactor in nearly 100 enzymes (B.L.Vallee *et al.*,1990). Zinc is required throughout the body to activate enzymes and to form metallo enzymes. These enzymes are required for metabolism and protein synthesis. The daily requirement of manganese for human is about 2-3 mg and zinc is 15 mg.

As salt manufacture is a semi agricultural operation, during monsoon season, the harvest would be stopped. During the study period heavy rain was experienced by all the salt works during the months of November and December 2013 and hence salt samples were not available during these two months and also in the following month January 2014.

II. STUDY AREA

Salt samples from the the two selected salt works, Kovalam and Puthalam (SW₁ and SW₂) of Kanyakumari District and the two selected salt works, Vepalodai and Sahupuram (SW₃ and SW₄) of Tuticorin District. The salt samples were collected and monthly for the period of one year from July 2013 to June 2014.

TABLE I THE MEAN AND STANDARD DEVIATION OF VARIOUS SALT QUALITY CONTENT (%) OF THE SALT SAMPLES DURING THE STUDY PERIOD JULY 2013 – JUNE 2014

| Salt Works | Moisture Content (%) | Insoluble Impurities (%) | CaSO ₄ Content (%) | MgSO ₄ Content (%) | MgCl ₂ Content (%) | NaCl Content (%) |
|------------|----------------------|--------------------------|-------------------------------|-------------------------------|-------------------------------|------------------|
| SW1 | 7.23±0.77 | 0.84±0.05 | 0.38±0.05 | 1.37±0.07 | 1.21±0.10 | 89.25±0.61 |
| SW2 | 9.26±0.41 | 0.93±0.38 | 0.24±0.03 | 1.54±0.06 | 1.48±0.19 | 86.51±0.71 |
| SW3 | 3.02±0.16 | 0.24±0.02 | 0.43±0.03 | 1.14±0.13 | 1.22±0.03 | 94.01±0.38 |
| SW4 | 5.11±0.62 | 2.15±0.41 | 0.88±0.10 | 1.40±0.16 | 1.81±0.22 | 85.90±1.03 |

The mean insoluble impurities presented in the samples under investigation had a wide range between 0.24 ± 0.02 to 2.15 ± 0.41 %. The maximum amount of insoluble impurities was present in SW₄ and the minimum amount was present in SW₃ throughout the period of investigation. The insoluble impurities in the salt works are due to the bad treatment of the pan floors and also the monsoonal rain, which spoil the salt pan floors at regular interval and also muddy area. In the present study, the insoluble impurities in the salt work SW₃, the value 0.24 ± 0.02% is almost equal to the value 0.1 to 1% reported for sea salt in other parts of India. In SW₃, the pond managements, pan flooring is once done properly.

Calcium sulphate is the most persistent companion of solar sea salt (Food and Nutrition Board, 2004). The mean value of calcium sulphate present in the samples was ranged from 0.24 ± 0.03 to 0.88 ± 0.10%. The minimum mean value of 0.24 ± 0.03% was estimated in the salt samples of SW₂. The maximum value of 0.88 ± 0.10% was estimated in the salt

III. MATERIALS AND METHODS

Salt samples were packed in air tight polythene containers. The different salt quality parameters like, the moisture content, the insoluble matter, calcium sulphate, magnesium sulphate, magnesium chloride and sodium chloride content were analyzed on wet basis as per the PFA ACT using standard methods. The determination of all the chemical parameters was done by the standard procedures (A.I.Vogel, 1961). The heavy metals of copper, iron, zinc, manganese, and chromium were analysed using Atomic Absorption Spectrophotometer (AAS). An average and standard deviation were taken of the data presented monthly during the study period.

IV. RESULTS AND DISCUSSION

Table I provides the mean and standard deviation of salt quality parameters. The mean value of moisture content of the salt samples fluctuated between 4.02±0.16 to 9.26 ± 0.41%, while samples had a maximum mean value of 9.26 ± 0.41% recorded in SW₂ and a minimum mean value of 3.02 ± 0.16 % recorded in SW₃. The fluctuations in the percentage of moisture clearly indicated that the atmospheric climate influenced the moisture content. The comparison between the salt samples from Kanyakumari and Tuticorin Districts, the moisture content was lower in Tuticorin District samples ie (SW₃ and SW₄).

samples of SW₄. Celtic sea salt had a mean calcium sulphate content of 0.435 %, whereas the earlier report (A.Chidambarathanu, 1997). This value is almost same in the present study. The mean value of calcium sulphate content 0.43% was estimated in SW₃.

The magnesium sulphate and magnesium chloride impurities in the sea salt were within the ratio 1:1:5(L.Crisman *et al.*, 1996). But the salts of the present study magnesium sulphate values are slightly greater or almost equal to the magnesium chloride content. The same observation had been noticed by earlier workers in the salt samples of the study area in Kanyakumari District ie.,(SW₁ & SW₂) where the magnesium sulphate content was higher than magnesium chloride (G.Leemarose,2007). In the present study the mean value of magnesium sulphate content was maximum 1.54 ± 0.06% in SW₂. The magnesium sulphate content was minimum 1.14 ± 0.13% in SW₃.

In the present study mean value of magnesium chloride content in the salt samples ranged from 1.21 ± 0.10 to $1.81 \pm 0.22\%$. The maximum chloride content $1.81 \pm 0.22\%$ was observed in the salt samples of SW₄. The minimum value of $1.21 \pm 0.10\%$ was observed in SW₁.

The accumulation of magnesium sulphate and magnesium chloride in the salt samples was mainly due to the non elimination of bittern in the crystallizer ponds after each scraping of salt. In SW₄, salt work had no proper channel for bittern. The magnesium rich liquor gets stagnated and remains in the crystallizers for a long time. Hence, when sodium chloride crystallizers, they trap more and magnesium salts in the lattice making it impurities.

SW₁ and SW₃ both salt works used pure sea brine, harvested salt of the maximum mean sodium chloride content of $94.01 \pm 0.38\%$ was observed in SW₃. The minimum mean sodium chloride content of $85.9 \pm 1.03\%$ was observed in SW₄. Because of the climatic condition (dry weather) and proper pond management, proper layout and elimination of bittern during the salt production process enhance the maximum mean sodium chloride content in SW₃ i.e., $94.01 \pm 0.38\%$. The mean sodium chloride content of $89.25 \pm 0.61\%$ in SW₁, was confirmed by earlier workers (A.Chidambarathanu, 1997). SW₂ and SW₄ both salt works

are used subsoil brine for salt production, harvested salt samples with mean sodium chloride content was $86.51 \pm 0.71\%$, $85.9 \pm 1.03\%$. The salt quality parameters revealed that moisture content was maximum during the rainy season while the sodium chloride content was maximum during the other seasons.

An increase in concentration of magnesium salts in sodium chloride crystals resulted in an increase in amount of moisture content resulting in reduction of sodium chloride content (J.G.Walmsley, 1997). The data on the monthly variation in the salt quality parameters revealed that moisture content was maximum during the rainy season while the sodium chloride content was maximum during other seasons. The salt samples having maximum moisture content in Kanyakumari salt works than Tuticorin District because of the climatic conditions.

Table II provides the mean and standard deviation of heavy metal concentration (ppm) of salt samples. The accumulation of copper in the salt harvested ranged from 0.011 ± 0.0049 to 0.079 ± 0.0053 ppm. Among these, the maximum of 0.079 ± 0.0053 ppm and the minimum of 0.011 ± 0.0049 ppm values were recorded at the salt samples from SW₄, SW₃ respectively.

TABLE II THE MEAN AND STANDARD DEVIATION OF HEAVY METAL CONCENTRATION (PPM) OF SALT SAMPLES DURING THE STUDY PERIOD JULY 2013 – JUNE 2014

| Salt Works | Copper(ppm) | Iron(ppm) | Zinc(ppm) | Manganese(ppm) | Chromium(ppm) |
|-----------------|--------------|---------------|--------------|----------------|---------------|
| SW ₁ | 0.028±0.0034 | 0.048±0.0037 | 0.049±0.0038 | 0.15±0.0235 | 2.5±0.2759 |
| SW ₂ | 0.072±0.0053 | 0.0316±0.0052 | 0.12±0.0208 | 0.19±0.0350 | 0.79±0.0660 |
| SW ₃ | 0.011±0.0049 | 0.046±0.0017 | 0.038±0.0034 | 0.07±0.0322 | 1.4±0.2698 |
| SW ₄ | 0.079±0.0029 | 0.028±0.0037 | 0.16±0.0255 | 0.20±0.0135 | 0.89±0.0899 |

The accumulation of iron in the salt harvested ranged from 0.028 ± 0.0037 to 0.048 ± 0.0037 ppm. Among these, the maximum 0.048 ± 0.0037 ppm at SW₁ and the minimum 0.028 ± 0.0037 ppm values were recorded at the salt samples from SW₄.

The accumulation of zinc in the salt harvested ranged from 0.038 ± 0.0034 to 0.16 ± 0.0255 ppm. Among these, the maximum 0.16 ± 0.0255 ppm at SW₄ and the minimum 0.038 ± 0.0034 ppm values were recorded at the salt samples from SW₃.

The accumulation of manganese in the salt harvested ranged from 0.07 ± 0.0322 to 0.20 ± 0.0135 ppm. Among these, the maximum 0.20 ± 0.0135 ppm at SW₄ and the minimum 0.07 ± 0.0322 ppm values were recorded at the salt samples from SW₃.

The accumulation of chromium in the salt harvested ranged from 0.79 ± 0.0660 to 2.5 ± 0.275 ppm. Among these, the maximum 2.5 ± 0.275 ppm at SW₁ and the minimum $0.79 \pm$

0.0660 ppm values were recorded at the salt samples from SW₂.

In the present study, salt sample obtained from subsoil brine (i.e., SW₂, SW₄) accumulated maximum copper content of 0.072 ppm and 0.079 ppm, whereas sea salt (SW₁, SW₃) accumulated the minimum copper content of 0.028 ppm and to 0.011 ppm. The mean copper content reported in the commercial sea salt is 0.0014 ppm, which is almost equal to the value recorded for the sea salt 0.011 ppm in the present investigation. The amount of zinc accumulated in the salt samples ranged from 0.049 to 0.16 ppm. According to the UK food standards committee report, zinc levels in food should not exceed 50 ppm. This shows the zinc content present in the salt samples is below the acceptable limit.

The low iron concentration of 0.020 ppm was reported in the sea waters of the Cadiz Bay (Jose Usero, 2003). In the salt samples, the iron content ranged from 0.028 to 0.048 ppm. The mean iron content recorded in the commercial sea salt was 0.015 ppm, which is slightly higher value to the minimum value reported in the present study.

When compared to other heavy metals, manganese content was present higher in salt samples. In the salt samples, the amount on manganese ranged from 0.07 to 0.20 ppm. The mean value of manganese reported in the commercial salt was 0.038 ppm.

Moreover, the amount of chromium recorded in the salt samples was ranging from 0.79 to 2.5 ppm. This value was agreeable with the earlier report (G. Leemarose, 2007). The amount of chromium in the salt sample was lesser than (3.5 ppm) the recommended by the west Australian food and Drug Regulations List.

V. CONCLUSION

The salt samples have maximum moisture content in Kanyakumari District than Tuticorin District because of the climatic conditions. The salt works used pure sea brine, harvested salt of the maximum sodium chloride content than sub-soil brine used salt works. Because of the climatic condition (dry weather) and proper pond management, proper layout and elimination of bittern during the salt production process enhance the quality of salt. In both the Districts, Kanyakumari and Tuticorin, Salt production using sea brine is having almost similar concentration of heavy metals, likewise salt production using sub soil brine.

REFERENCES

- [1] Earl Friden, "The Chemical Elements of Life", *Scientific American Journal*, 1972.
- [2] D. W. Kaufmann, Sodium chloride, ACS Monograph 145, Reinhold publishing Corp., New York, 1960.
- [3] United Nations Industrial Development Organisation, Vienna, Proc. Of Expert Group meeting, Rome, September, pp. 25-29, 1968.
- [4] S. C. Agarwal, *The Salt Industry India*, The Manger of publications, Government Press of India, Delhi, pp. 647, 1956.
- [5] P.Sorgeloos and W. Tackaert, *Proc.of the Intl.symp. on Biotechnology of salt pond*, Tang Gu, PR China pp.13-21,1990.
- [6] G. Leema Rose, Ph.D. Thesis, M. S. University, Tirunelveli, 2007.
- [7] Rainbow, The significance of trace metal concentration in decapods. *Symp. Zool. Soc. London*, Vol. 59, pp. 291 – 313, 1986.
- [8] P. Miramand, T. Guyot, H. Rybarczy, B. Elkaim, P. Mouny, J. C. Dauvin and C. Bessineton, *Eusteries*, Vol. 24, No. 66, pp. 1056-1065, 2001.
- [9] B. L. Vallee and D. S. Auld, *Biochemistry*, Vol. 29, pp. 5647-5659, 1990.
- [10] A. I. Vogel, A Text book of Quantitative Inorganic Analysis, 3rd Edn., ELBS and Longmans, London, 1961.
- [11] Food and Nutrition Board, Institute of Medicine. Sodium and chloride Dietary Reference Intakes for water, potassium, sodium, chloride and sulphate, Washington, D.C: National academic Press, pp. 247 – 392, 2004.
- [12] A. Chidambarathanu, Ph. D. Thesis, M. S. University, Tirunelveli, 1997.
- [13] L. Crisman, L. J. Chapman and C.A.Chapman, *Geotimes*, pp. 23-23, July, 1996.
- [14] C.Vaithyanathan Ph.D Thesis, M.S.University, Tirunelveli, 2004.
- [15] J. G. Walmsley, Mediterranean saline distribution, salt production and conservation. In: *Proceedings on Nature and Workmanship, Artificial wetlands in the Mediterranean Coast*, INSULA, UNESCO Building, Paris,pp. 152, 1997.
- [16] Jose Usero, Carmen Izquierdo, Jose Morillo and Ignacio Gracia, *Environment International*, Vol. 29, pp. 949-956, 2003.